BUD COMPLEX


FERTILISER - A WATER SOLUBLE FERTILISER FOR GRAPES AND TREE CROPS

3.9%

MADE IN FRANCE

Agronutrition

Precision technologies



Specifications

PACK SIZE 10 KG

2.4%

FORMULATION

SHELF LIFE OF THE PRODUCT 24 months APPLICATION Foliar

FORM Powder (SP) DENSITY

STORAGE CONDITIONS Store away from humidity.

Specific Mineral balance

Agronomic interests

ROLE OF NUTRIENT(S)	PRODUCT SPECIFICATIONS	FEATURES AND BENEFITS OF THE FORMULATION
Phosphorus: Energy carrying/Membrane constituents/ Protein synthesis	A soluble micronutrient blend tailored to meet the typical nutrient requirements of all fruit trees after leaf fall.	The unique formulation of this product guarantees maximum foliar uptake, without the risk of phytotoxicity, designed to achieve nutrient absorption in less than 6 hours.
Magnesium: Photosynthesis/Synthesis of carbohydrates and lipids	BUD COMPLEX contains a balanced ratio of fully and plant-available nutrients formulated for easy and straight forward mixing and application.	This product is formulated with rigorously selected raw materials, guaranteed by quality control, to ensure a perfect safety of use and to optimize the absorption
Boron: Role on flowering and fruit set/Migration on sugars/ Protein synthesis/Meristematic activity and growth/ Use of auxin in the plant	BUD COMPLEX is highly concentrated in elements preventing and thus efficiently correcting the deficiency in Boron, Zinc, Magnesium and Phosphorus.	through the leaves.
Zinc: Growth hormone synthesis/Enzymatic functioning/Protein synthesis	The rigorous selection of the raw materials guarantees it to be 100% soluble in the treatment mix.	

Directions of use

CROP RECOMMENDATIONS

> GRAPES

Rate: 4-6 kg/ha in 500-1000 L of clean water. Timing: apply after each harvest before leaf fall, or when sufficient green leaf is present during the spring leaf development period.

> TREE CROPS

Rate: 6 kg/ha in 500-1000 L of clean water. Timing: apply after each harvest before leaf fall.

Instructions for use

For best results, use sufficient water to achieve an even crop coverage to the point of runoff.

Avoid spraying in very strong sunlight and/or very high temperatures. If possible, apply during evening or early morning especially if day-time temperatures exceed 28°C. Very young foliage and crops under stress may be more susceptible to scorch. Lower doses should be applied in these circumstances and the application interval reduced accordingly. Refer to the Crop Recommendations table for application rates and timings for individual crops.

Precautions

MIXING

II. Fill the spray tank with half the required amount of water and start agitation. 2. Gradually add the required amount of BUD COMPLEX and then add the rest of the water, maintaining agitation continuously. When combining with other components in a tank-mix, always add BUD COMPLEX last. Do not allow the mixture to stand without agitation. 3. After spraying, clean and rinse the spraying equipment thoroughly.

IMPORTANT: respect the uses, doses, conditions and instructions for use mentioned on the packaging, which are determined according to the characteristics of the product and the applications for which it is recommended. On this basis, conduct the crop and treatments according to the best agricultural practices taking into account, under your responsibility, all the specific factors concerning your farm, such as soil characteristics, weather conditions, cropping methods, cultivars and their specific resistances. AGRONUTRITION guarantees the quality its products only in their initial packaging over the period indicated on the manufacturer's certificates. They guarantee their conformity to the composition indicated on the packaging and to the regulations currently in force. Precautionary measure brochures are available on request. To request a MSDS, thank you to contact: infoau@desangosse.com

