

CIGOPHOL K

FERTILISER - K AND P FERTILISER SOLUTION WITH BORON

Precision
technologies

Agro nutrition

N	P	K	Mg	S	Ca
	105	264			
B	Cu	Fe	Mn	Mo	Zn
8					



In g/L for liquid products

Specifications

PACK SIZE

10 L

SHELF LIFE OF THE PRODUCT

24 months

FORM

Liquid (SL)

STORAGE CONDITIONS

Store away from frost.

FORMULATION

Specific Mineral Balance

APPLICATION

Foliar

DENSITY

1.46

Agronomic interests

ROLE OF NUTRIENT(S)

Phosphorus :

- Energy carrying
- Membrane constituents
- Protein synthesis

Potassium :

- Synthesis and transfer of sugars
- Photosynthesis
- Water regulation and management

PRODUCT SPECIFICATIONS

Cigophol K is specifically designed and formulated to optimise maturity and quality of the crop at harvest.

Through foliar uptake, it helps with the transfer of assimilates from the leaves to the fruit.

Cigophol K is a 100% soluble Potassium concentrate formulated to provide maximum effect in synergy with phosphorus.

FEATURES AND BENEFITS OF THE FORMULATION

The unique formulation of this product guarantees maximum foliar uptake, without the risk of phytotoxicity, designed to achieve nutrient absorption in less than 6 hours.

This product is formulated with rigorously selected raw materials, guaranteed by quality control, to ensure a perfect safety of use and to optimise the absorption through the leaves.

Directions of use

CROP RECOMMENDATIONS

- > **GRAPES** - Rate: 3-5 L/ha in 200 L minimum of water - Timing: 2-4 applications applied at 10-15 day intervals starting from fruit set and during ripening.
- > **FRUIT TREES AND KIWIFRUIT** - Rate: 5 L/ha in 200 L minimum of water - Timing: 3 applications applied at 10-15 day intervals starting at the beginning of fruit growth and up to 2 weeks before harvest.
- > **FIELD CROPS** - Rate: 3-5 L/ha in 200 L minimum of water - Timing: 3 to 5 applications on sufficiently developed foliage.
- > **VEGETABLES** - Rate: 3-5 L/ha in 500 L minimum of water - Timing: 3 to 5 applications on sufficiently developed foliage.

Instructions for use

For best results, use sufficient water to achieve an even crop coverage to the point of run-off. Avoid spraying in very strong sunlight and/or very high temperatures. If possible, apply during evening or early morning especially if day-time temperatures exceed 28°C. Very young foliage and crops under stress may be more susceptible to scorch. Lower doses should be applied in these circumstances and the application interval reduced accordingly. Refer to the Crop Recommendations table for application rates and timings for individual crops. Use outdoors only.

Precautions

MIXING

1. Fill the spray tank with half the required amount of water and start agitation.
2. Add the required amount of Cigophol K and then add the rest of the water, maintaining agitation continuously. When combining with other components in a tank-mix, always add Cigophol K last. Do not allow the mixture to stand without agitation.
3. After spraying, clean and rinse the spraying equipment thoroughly.

IMPORTANT: respect the uses, doses, conditions and instructions for use mentioned on the packaging, which are determined according to the characteristics of the product and the applications for which it is recommended. On this basis, conduct the crop and treatments according to the best agricultural practices taking into account, under your responsibility, all the specific factors concerning your farm, such as soil characteristics, weather conditions, cropping methods, cultivars and their specific resistances. AGRONUTRITION guarantees the quality its products only in their initial packaging over the period indicated on the manufacturer's certificates. They guarantee their conformity to the composition indicated on the packaging and to the regulations currently in force. Precautionary measure brochures are available on request. To request a MSDS, thank you to contact: infoau@desangosse.com