

# FERFOL MB

FERTILISER - IRON CHELATE EDTA

Precision  
technologies



N	P	K	Mg	S	Ca
100			18		
B	Cu	Fe	Mn	Mo	Zn
10		35			



In g/L for liquid products

## Specifications

### PACK SIZE

10 L

### SHELF LIFE OF THE PRODUCT

24 months

### FORM

Liquid (SL)

### STORAGE CONDITIONS

Store away from frost and keep in original container out of direct sunlight and tightly closed.

### FORMULATION

Specific Mineral balance in foliar

### APPLICATION

Foliar

## Agronomic interests

### ROLE OF NUTRIENT(S)

**Boron :**  
Role on flowering and fruit set / Migration on sugars / Protein synthesis / Meristematic activity and growth / Use of auxin in the plant

**Iron :**  
Transpiration / Chlorophyll formation / Functioning of the photosynthetic systems / Protein metabolism / Nitrate reduction

**Magnesium :**  
Photosynthesis / Synthesis of carbohydrates and lipids

### PRODUCT SPECIFICATIONS

FERFOL MB contains chelated Iron (EDTA). The EDTA chelated form acts fast and its systemic action optimises photosynthetic activity in the plant.

FERFOL MB is a ready to use liquid product, safe to use and handle.  
The synergy between Iron, Boron and Magnesium elements increases the efficiency of the treatment.

FERFOL MB is a polyvalent product which is suitable to numerous crops. It improves homogeneity and quality of harvested crops.

### FEATURES AND BENEFITS OF THE FORMULATION

The unique formulation of this product guarantees maximum foliar uptake, without risk of phytotoxicity. Designed to achieve nutrient absorption in less than 6 hours.

This product is formulated with rigorously selected raw materials, guaranteed by quality control, to ensure a perfect safety of use and to optimise the absorption through the leaves.

## Directions of use

### CROP RECOMMENDATIONS

#### FOLIAR APPLICATION

Maximum concentration : 1% (1L in 100L of water)

#### > GRAPES

Rate: 3 L/ha – Timing: Separated bunches - start flowering - fruit set - bunch closure

#### > FRUIT TREES (stone and pip fruits)

Rate: 5 L/ha – Timing: start flowering, petals fall, fruit set and young fruit

> **FIELD CROPS AND VEGETABLES** - Rate: 5L/ha – Timing: 2-3 applications from 10 days interval on well developed foliage

### Instructions for use

For best results, use sufficient water to achieve an even crop coverage to the point of run-off. Avoid spraying in very strong sunlight and/or very high temperatures. If possible, apply during evening or early morning especially if day-time temperatures exceed 25°C. Very young foliage and crops under stress may be more susceptible to scorch. Lower doses should be applied in these circumstances and the application interval reduced accordingly. Refer to the Crop Recommendations table for application rates and timings for individual crops. Use outdoors only. Fill the spray tank with half the required amount of water and start agitation. Shake container well and add the required amount of our product, then add the rest of the water maintaining agitation continuously. After spraying, clean and rinse the spraying equipment thoroughly.

## Precautions

### COMPATIBILITY:

Always read the product labels and follow the manufacturers' instructions for all products at all times. Some settings, outside the control of the manufacturer or distributor, may have repercussions on the performance of co-applied products. Therefore, the co-application is made at the risk of the end user. When combining with other components in a tank-mix, always add this product last. If there is any doubt at all, consult the manufacturer or distributor concerned.

**IMPORTANT :** Respect the uses, doses, conditions and instructions for use mentioned on the packaging, which are determined according to the characteristics of the product and the applications for which it is recommended. On this basis, conduct the crop and treatments according to the best agricultural practices taking into account, under your responsibility, all the specific factors concerning your farm, such as soil characteristics, weather conditions, cropping methods, cultivars and their specific resistances. AGRONUTRITION guarantees the quality its products only in their initial packaging over the period indicated on the manufacturer's certificates. They guarantee their conformity to the composition indicated on the packaging and to the regulations currently in force. Precautionary measure brochures are available on request. **To request a MSDS, thank you to contact : infoau@desangosse.com**